

Improving fluid intelligence with training on working memory

Susanne M. Jaeggi*^{†‡}, Martin Buschkuhl*^{†‡}, John Jonides*, and Walter J. Perrig[†]

*Department of Psychology, University of Michigan, East Hall, 530 Church Street, Ann Arbor, MI 48109-1043; and [†]Department of Psychology, University of Bern, Muesmattstrasse 45, 3012 Bern, Switzerland

Edited by Edward E. Smith, Columbia University, New York, NY, and approved March 18, 2008 (received for review February 7, 2008)

Fluid intelligence (*Gf*) refers to the ability to reason and to solve new problems independently of previously acquired knowledge. *Gf* is critical for a wide variety of cognitive tasks, and it is considered one of the most important factors in learning. Moreover, *Gf* is closely related to professional and educational success, especially in complex and demanding environments. Although performance on tests of *Gf* can be improved through direct practice on the tests themselves, there is no evidence that training on any other regimen yields increased *Gf* in adults. Furthermore, there is a long history of research into cognitive training showing that, although performance on trained tasks can increase dramatically, transfer of this learning to other tasks remains poor. Here, we present evidence for transfer from training on a demanding working memory task to measures of *Gf*. This transfer results even though the trained task is entirely different from the intelligence test itself. Furthermore, we demonstrate that the extent of gain in intelligence critically depends on the amount of training: the more training, the more improvement in *Gf*. That is, the training effect is dosage-dependent. Thus, in contrast to many previous studies, we conclude that it is possible to improve *Gf* without practicing the testing tasks themselves, opening a wide range of applications.

cognitive training | transfer | individual differences | executive processes | control processes

Fluid intelligence (*Gf*) is a complex human ability that allows us to adapt our thinking to a new cognitive problem or situation (1). *Gf* is critical for a wide variety of cognitive tasks (2), and it is considered one of the most important factors in learning. Moreover, *Gf* is closely related to professional and educational success (3–6), especially in complex and demanding environments (7). There is considerable agreement that *Gf* is robust against influences of education and socialization, and it is commonly seen as having a strong hereditary component (2, 8, 9). *Gf* can be compromised as seen in the effects of certain manipulations that threaten one's group membership (10). But can *Gf* be improved by any means?

In the domain of psychopharmacology, although there is a market for so-called “smart drugs,” there is no study showing evidence for a drug-related increase in *Gf* in healthy adults although there are certain psychomotor stimulants and D2 dopamine-receptor agonists that have effects on isolated cognitive processes (11, 12). Beyond the psychopharmacology, there is a growing interest in whether computer and video games may increase IQ. But in contrast to suggestive advertisements, there is no empirical evidence that computer games enhance anything beyond task-specific performance (13, 14) and selective visuospatial attention (15).

Of course, one can easily increase performance in tests of *Gf* by simply practicing the tests themselves (16). However, it has been demonstrated that practice on these tests decreases their novelty and with that the underlying *Gf*-processes (5) so that the predictive value of the tests for other tasks disappears (17). These findings are compatible with a long history of research on cognitive training in psychological and educational science showing that, although performance on trained tasks can increase

dramatically, transfer of this learning to other tasks or domains remains shockingly rare (18–21).

Despite the many failures to find transfer in any domain, the sheer importance of identifying tasks that can lead to improvement in other tasks recommends continued investigation of transfer effects. With respect to *Gf*, the issue is whether one can identify a task that shares many of the features and processes of *Gf* tasks, but that is still different enough from the *Gf* tasks themselves to avoid mere practice effects. A recently proposed hypothesis by Halford *et al.* (22) might serve as a useful framework for the design of a transfer study in which one would like to improve *Gf* by means of a working memory task. Their claim is that working memory and intelligence share a common capacity constraint. This capacity constraint can be expressed either by the number of items that can be held in working memory or by the number of interrelationships among elements in a reasoning task. The reason for a common capacity limitation is assumed to lie in the common demand for attention when temporary binding processes are taking place to form representations in reasoning tasks (22). Other authors came to a related conclusion, stating that *Gf* and working memory are primarily related through attentional control processes (23, 24). Furthermore, Carpenter *et al.* (1) have proposed that the ability to derive abstract relations and to maintain a large set of possible goals in working memory accounts for individual differences in typical tasks that measure *Gf*.

The underlying neural circuitries provide additional evidence for the shared variance between working memory and *Gf* in that both seem to rely on similar neural networks, most consistently located in lateral prefrontal and parietal cortices (23, 25). Therefore, it seems plausible that the training of a certain neural circuit might lead to transfer on other tasks that engage similar or at least overlapping neural circuits.

Although working memory capacity and *Gf* may share common variance, they are far from being isomorphic (26, 27). That is, there are factors other than working memory capacity contributing to individual differences in *Gf*. Nevertheless, we propose that, with a training intervention that strongly relies on binding processes and attentional control, it may be possible to produce transfer effects from a trained task to a reasoning task in which performance relies to a large extent on the same processes. There are, indeed, some studies showing that training on working memory with young healthy adults may lead to effects that go beyond a specific training effect (28, 29). So, it seems that there is some potential for transfer after training on working memory.

To investigate whether training on working memory leads to transfer to *Gf*, we conducted four individual experiments all

Author contributions: S.M.J. and M.B. contributed equally to this work; and S.M.J., M.B., J.J., and W.J.P. wrote the paper.

The authors declare no conflict of interest.

This article is a PNAS Direct Submission.

[†]To whom correspondence may be addressed. E-mail: sjaeggi@umich.edu or mbu@umich.edu.

© 2008 by The National Academy of Sciences of the USA




Fig. 1. The n -back task that was used as the training task, illustrated for a 2-back condition. The letters were presented auditorily at the same rate as the spatial material was presented visually.

using a newly developed training paradigm consisting of a very demanding working memory task, illustrated in Fig. 1. In this task, participants saw two series of stimuli that were synchronously presented at the rate of 3 s per stimulus. One string of stimuli consisted of single letters whereas the other consisted of individual spatial locations marked on a screen. The task was to decide for each string whether the current stimulus matched the one that was presented n items back in the series. The value of n varied from one block of trials to another, with adjustments made continuously for each participant based on performance. As performance improved, n incremented by one item; as it worsened, n decremented by one item. Thus, the task changed adaptively so that it always remained demanding, and this demand was tailored to individual participants. This form of training engaged processes required for the management of two simultaneous tasks; it engaged executive processes required for each task; and it discouraged the development of task-specific strategies and the engagement of automatic processes because of the variation in n and because of the inclusion of two different classes of stimuli.

The aim of the training intervention was the investigation of the effects of training on the working memory task and its impact on Gf . We accomplished this investigation by pretesting participants on a measure of Gf and then posttesting them on another form of this measure. Because we hypothesized that any alteration of the processing system would take some time to be effective, an important difference among the four experiments was the number of training sessions between pre- and posttests, ranging from 8 to 19 sessions. To control for mere retest effects, the performance of the trained groups was compared with control groups who were also assessed on Gf , but who were not trained between the two testing sessions.

Results

Analyses of the training functions revealed that all four training groups improved in their performance on the working memory task comparably (Fig. 2).[§] What interests us most, however, is the dramatic improvement on the test of Gf in the trained groups (Fig. 3a). Although the gain in the control groups was also significant, presumably because of retest effects ($t(34) = 2.08$;

[§]Note that the specific training gain was well explained by a linear function for all groups (8 days training: $R^2 = 0.78$, $F(1,134) = 473.44$, $P < 0.001$; 12 days training: $R^2 = 0.81$, $F(1,202) = 863.59$, $P < 0.001$; 17 days training: $R^2 = 0.73$, $F(1,287) = 769.63$; 19 days training: $R^2 = 0.79$, $F(1,321) = 1230.23$, $P < 0.001$).




Fig. 2. Performance increase in the trained task shown separately for each training group. For each session, the mean level of n achieved by the participants is presented. The level of n depends on the participants' performance.

$P < 0.05$; Cohen's $d = 0.25$), the improvement in the groups that received the apparent benefit of training was substantially superior ($t(33) = 5.53$; $P < 0.001$; Cohen's $d = 0.65$), which was confirmed by the significant group \times test-session interaction ($F(1,67) = 5.27$; $P < 0.05$; $\eta_p^2 = 0.07$). A subsequent analysis of the gain scores (posttest minus pretest) as a function of training time (8, 12, 17, or 19 days) showed that transfer to fluid intelligence varied as a function of training time ($F(3,30) = 9.25$; $P < 0.001$; $\eta_p^2 = 0.48$; Fig. 3b). Analyses of covariance (ANCOVA) with the factor group (trained vs. control), the posttest scores as the dependent variable, and the pretest scores as the covariate revealed a trend for group differences after 12 days ($F(1,19) = 1.93$; $P = 0.09$; $\eta_p^2 = 0.09$), and statistically significant group differences after 17 ($F(1,13) = 4.65$; $P < 0.05$; $\eta_p^2 = 0.26$), and 19 training days ($F(1,12) = 4.53$; $P < 0.05$; $\eta_p^2 = 0.27$). Post hoc analyses (Gabriel's procedure; two-tailed) for the training group revealed significant differences between the following groups: 8 vs. 17 days ($P < 0.01$); 8 vs. 19 days ($P < 0.001$); and 12 vs. 19 days ($P < 0.01$). There was a trend for a difference between 12 and 17 days ($P = 0.06$). These analyses indicate that the gain in fluid intelligence was responsive to the dosage of training.

It is important to note that the gain in Gf is strictly training-related and not due to preexisting individual differences in intelligence or working memory. There was an effect of training ($F(1,68) = 6.38$; $P < 0.05$; $\eta_p^2 = 0.09$) irrespective of initial Gf as shown by dividing the sample into high and low performers in Gf at pretest (by median split). However, there was also a main effect of performance group ($F(1,68) = 4.56$; $P < 0.05$; $\eta_p^2 = 0.07$), showing that participants with initially lower Gf generally showed even larger gains in Gf . There was no significant performance group by training-gain interaction ($F < 1$).

Further, the gain in Gf was not dependent on initial working memory capacity as assessed either by pretest performance in a digit-span task ($F(1,68) < 0.3$) or a reading-span task ($F(1,52) < 0.1$; note that reading span was not assessed in the 8-day group). Therefore, our cognitive training proved to be useful for all participants and had no adverse effects for participants with high initial working memory capacity.

Additional analyses showed that there was training-related transfer on the digit-span task (group \times session interaction: $F(1,67) = 14.08$; $P < 0.001$; $\eta_p^2 = 0.17$), but not on the reading span task, in which both groups improved equally in the posttest

skills not measured by simpler working memory tasks that are engaged by our training task and that train skills needed in measures of *Gf*. It may still be the case that training on the dual *n*-back task promotes development of these non-capacity skills. A line of evidence consistent with this idea shows that individual differences in both working memory span and in *n*-back tasks are related to individual differences in *Gf* (23, 25, 32).

The finding that the transfer to *Gf* remained even after taking the specific training effect into account seems to be counterintuitive, especially because the specific training effect is also related to training time. The reason for this capacity might be that participants with a very high level of *n* at the end of the training period may have developed very task specific strategies, which obviously boosts *n*-back performance, but may prevent transfer because these strategies remain too task-specific (5, 20). The averaged *n*-back level in the last session is therefore not critical to predicting a gain in *Gf*; rather, it seems that working at the capacity limit promotes transfer to *Gf*.

Of particular importance is the finding that preexisting inter-individual differences in *Gf* as measured in the pretest are not related to the training-related gain in *Gf*. This finding indicates that the effect of training is not restricted to participants within a certain range of cognitive abilities. Both initial low-*Gf* as well as initial high-*Gf* participants profit from training similarly. Still, although the interaction was not reliable, we remain cautious about this result because numerically the low-*Gf* participants showed somewhat larger gains than the high-*Gf* participants. Of course, this result may be accounted for by regression to the mean, but it may also be that the training was of truly greater benefit to lower *Gf* participants, if not reliably so in our study.

The dose-responsive gain in *Gf* indicates that the training benefit is not a threshold phenomenon. The constraints of our experiments do not permit us to know how much longer we could have continued training before failing to realize any further gains in *Gf*. This dose responsiveness is an important issue for further study because the exact plot of gain with training could have important practical implications for those interested in training fluid intelligence. Our study also does not permit us to know how long the training gain persists; longitudinal studies will be required to address that issue.

These limitations notwithstanding, our findings are of general significance because they provide evidence for enhancement of fluid intelligence by cognitive training different from training the test itself. The finding that cognitive training can improve *Gf* is a landmark result because this form of intelligence has been claimed to be largely immutable. Instead of regarding *Gf* as an immutable trait, our data provide evidence that, with appropriate training, there is potential to improve *Gf*. Moreover, we provide evidence that the amount of *Gf*-gain critically depends on the amount of training time. Considering the fundamental importance of *Gf* in everyday life and its predictive power for a large variety of intellectual tasks and professional success, we believe that our findings may be highly relevant to applications in education.

Materials and Methods

Participants and Procedure. For this study, we conducted four individual experiments involving a total of 70 healthy young participants (36 female; mean age, 25.6 years of age; SD, 3.3) recruited from the University of Bern community, 35 of whom performed working-memory training in four differ-

ent training settings (one of the participants failed to complete the required training sessions and was thus discarded from the data analysis, resulting in a final *N* of 34). These training groups were matched to four control groups who did not have training (*n* = 35). The crucial difference among the four training settings was the number of training sessions between pre- and posttests, ranging from 8 to 19 sessions (i.e., 8 days (*n* = 16), 12 days (*n* = 22), 17 days (*n* = 16), and 19 days (*n* = 15)), with the control groups receiving the pre- and posttesting at the same intervals as the trained groups. In each training setting, participants trained daily, except for the weekends. The posttest took place at least 1 day after the last training session, with the largest interval being 2 days.

Materials. Training task. For the training task, we used the same material as described by Jaeggi *et al.* (33), which was a dual *n*-back task where squares at eight different locations were presented sequentially on a computer screen at a rate of 3 s (stimulus length, 500 ms; interstimulus interval, 2,500 ms). Simultaneously with the presentation of the squares, one of eight consonants was presented sequentially through headphones. A response was required whenever one of the presented stimuli matched the one presented *n* positions back in the sequence. The value of *n* was the same for both streams of stimuli. There were six auditory and six visual targets per block (four appearing in only one modality, and two appearing in both modalities simultaneously), and their positions were determined randomly. Participants made responses manually by pressing on the letter "A" of a standard keyboard with their left index finger for visual targets, and on the letter "L" with their right index finger for auditory targets. No responses were required for non-targets.

In this task, the level of difficulty was varied by changing the level of *n* (34), which we used to track the participants' performance. After each block, the participants' individual performance was analyzed, and in the following block, the level of *n* was adapted accordingly: If the participant made fewer than three mistakes per modality, the level of *n* increased by 1. It was decreased by 1 if more than five mistakes were made, and in all other cases, *n* remained unchanged.

One training session comprised 20 blocks consisting of 20 + *n* trials resulting in a daily training time of ≈25 min.

Transfer tasks. We used standardized fluid intelligence tests, consisting of visual analogy problems of increasing difficulty. Each problem presents a matrix of patterns in which one pattern is missing. The task is to select the missing pattern among a set of given response alternatives. For the experiment with eight training sessions, we used the Raven's Advanced Progressive Matrices (RAPM) test, set II (35), whereas for all other experiments, we used the short version of the Bochumer Matrizen-Test (BOMAT) (36), a more difficult variant of the RAPM. For the RAPM, we used parallel forms for the pre- and posttesting by dividing the test into even and odd items (24); for the BOMAT, we used the published A and B versions. To keep the pre- and posttest sessions short enough, we allowed limited time (10 min) to complete the task, and the number of correct solutions provided in that time served as the dependent variable.[†]

To control for the impact of individual differences and gain in working memory capacity, a digit-span task (38), as well as a reading span task (39), was used in the pre- and postsession. However, the reading span task was not assessed in the 8-day group.

[†]Although this procedure differs from the standardized procedure, there is evidence that this timed procedure has little influence on relative standings in these tests, in that the correlation of speeded and non-speeded versions is very high (*r* = 0.95; ref. 37).

ACKNOWLEDGMENTS. We thank Daniela Blaser and Oliver Markes for help with data collection and the participants for their time and effort. The preparation of this manuscript was supported by Swiss National Science Foundation Fellowships PA001-117473 (to S.M.J.) and PBBE1-117527 (to M.B.) and by grants from the National Science Foundation and the National Institute of Mental Health (to J.J.).

1. Carpenter PA, Just MA, Shell P (1990) What one intelligence test measures: A theoretical account of the processing in the Raven Progressive Matrices Test. *Psychol Rev* 97:404–431.
2. Gray JR, Thompson PM (2004) Neurobiology of intelligence: Science and ethics. *Nat Rev Neurosci* 5:471–482.
3. Neisser U, *et al.* (1996) Intelligence: Knowns and unknowns. *Am Psychol* 51:77–101.
4. Rohde TE, Thompson LA (2007) Predicting academic achievement with cognitive ability. *Intelligence* 35:83–92.

5. te Nijenhuis J, van Vianen AEM, van der Flier H (2007) Score gains on g-loaded tests: No g. *Intelligence* 35:283–300.
6. Deary IJ, Strand S, Smith P, Fernandes C (2007) Intelligence and educational achievement. *Intelligence* 35:13–21.
7. Gottfredson LS (1997) Why g matters: The complexity of everyday life. *Intelligence* 24:79–132.
8. Cattell RB (1963) Theory of fluid and crystallized intelligence: A critical experiment. *J Educ Psychol* 54:1–22.

9. Baltes PB, Staudinger UM, Lindenberger U (1999) Lifespan psychology: Theory and application to intellectual functioning. *Annu Rev Psychol* 50:471–507.
10. Steele CM, Aronson JA (2004) Stereotype threat does not live by Steele and Aronson (1995) alone. *Am Psychol* 59:47–48; discussion 48–49.
11. Elliott R, et al. (1997) Effects of methylphenidate on spatial working memory and planning in healthy young adults. *Psychopharmacology* 131:196–206.
12. Kimberg DY, D'Esposito M, Farah MJ (1997) Effects of bromocriptine on human subjects depend on working memory capacity. *NeuroReport* 8:3581–3585.
13. Haier RJ, et al. (1992) Regional glucose metabolic changes after learning a complex visuospatial/motor task: A positron emission tomographic study. *Brain Res* 570:134–143.
14. Smith ME, McEvoy LK, Gevins A (1999) Neurophysiological indices of strategy development and skill acquisition. *Brain Res Cognit Brain Res* 7:389–404.
15. Green CS, Bavelier D (2003) Action video game modifies visual selective attention. *Nature* 423:534–537.
16. Bors DA, Vigneau F (2003) The effect of practice on Raven's Advanced Progressive Matrices. *Learn Individ Differ* 13:291–312.
17. Ackerman PL (1987) Individual differences in skill learning: An integration of psychometric and information processing perspectives. *Psychol Bull* 102:3–27.
18. Healy AF, Wohldmann EL, Sutton EM, Bourne LE, Jr (2006) Specificity effects in training and transfer of speeded responses. *J Exp Psychol Learn Mem Cognit* 32:534–546.
19. Singley MK, Anderson JR (1989) *The transfer of cognitive skill* (Harvard Univ Press, Cambridge, MA).
20. Ericsson AK, Delaney PF (1998) in *Working Memory and Thinking*, eds Logie R, Gilhooly KJ (Erlbaum, Hillsdale, NJ), pp 93–114.
21. Chase WG, Ericsson KA (1981) Skilled memory. *Cognitive Skills and Their Acquisition*, ed Anderson, J. R. (Lawrence Erlbaum Associates, Hillsdale, NJ), pp 141–189.
22. Halford GS, Cowan N, Andrews G (2007) Separating cognitive capacity from knowledge: A new hypothesis. *Trends Cognit Sci* 11:236–242.
23. Gray JR, Chabris CF, Braver TS (2003) Neural mechanisms of general fluid intelligence. *Nat Neurosci* 6:316–322.
24. Kane MJ, et al. (2004) The generality of working memory capacity: A latent-variable approach to verbal and visuospatial memory span and reasoning. *J Exp Psychol Gen* 133:189–217.
25. Kane MJ, Engle RW (2002) The role of prefrontal cortex in working-memory capacity, executive attention, and general fluid intelligence: An individual-differences perspective. *Psychon Bull Rev* 9:637–671.
26. Ackerman PL, Beier ME, Boyle MO (2005) Working memory and intelligence: The same or different constructs? *Psychol Bull* 131:30–60.
27. Conway AR, Kane MJ, Engle RW (2003) Working memory capacity and its relation to general intelligence. *Trends Cognit Sci* 7:547–552.
28. Olesen PJ, Westerberg H, Klingberg T (2004) Increased prefrontal and parietal activity after training of working memory. *Nat Neurosci* 7:75–79.
29. Westerberg H, Klingberg T (2007) Changes in cortical activity after training of working memory: A single-subject analysis. *Physiol Behav* 92:186–192.
30. Smith EE, Jonides J (1998) Neuroimaging analyses of human working memory. *Proc Natl Acad Sci USA* 95:12061–12068.
31. Oberauer K (2005) Binding and inhibition in working memory: Individual and age differences in short-term recognition. *J Exp Psychol Gen* 134:368–387.
32. Kane MJ, Conway AR, Miura TK, Colflesh GJ (2007) Working memory, attention control, and the N-back task: A question of construct validity. *J Exp Psychol Learn Mem Cognit* 33:615–622.
33. Jaeggi SM, et al. (2007) On how high performers keep cool brains in situations of cognitive overload. *Cognit Affect Behav Neurosci* 7:75–89.
34. Jonides J, et al. (1997) Verbal working memory load affects regional brain activation as measured by PET. *J Cognit Neurosci* 9:462–475.
35. Raven JC (1990) *Advanced Progressive Matrices: Sets I, II* (Oxford Univ Press, Oxford).
36. Hossiep R, Turck D, Hasella M (1999) *Bochumer Matrizenstest: BOMAT-Advanced-Short Version* (Hogrefe, Göttingen).
37. Frearson W, Eysenck HJ (1986) Intelligence, reaction time (RT), and a new odd-man-out RT paradigm. *Pers Individ Differ* 7:807–818.
38. Tewes U (1991) *Hamburg-Wechsler-Intelligenztest für Erwachsene (HAWIE-R)* (Hans Huber, Bern, Switzerland), rev 1991.
39. Daneman M, Carpenter PA (1980) Individual differences in working memory and reading. *J Verbal Learn Verbal Behav* 19:450–466.